

Prospects of observing pulsed radiation from gamma-ray pulsars with H.E.S.S.

O. C. de Jager¹, A. Konopelko², B.C. Raubenheimer¹ and B. Visser¹

¹ *Unit for Space Physics, Potchefstroom University, Potchefstroom, 2520, South Africa*

² *Max Planck Institut für Kernphysik, Postfach 103980, D-69029 Heidelberg, Germany*

Abstract. Observations and theoretical studies have demonstrated that the pulsed spectra of all gamma-ray pulsars terminate at energies below a few hundred GeV. In most cases we expect this cutoff energy E_o to be around 10 GeV. Only with next-generation ground-based γ -ray telescopes, which are expected to have non-zero trigger probabilities near 10 GeV, can we expect to detect pulsations. The large γ -ray fluxes below E_o , together with the associated sharp pulse profiles, compensate for the lack of imaging capability near threshold. For H.E.S.S. we find that the pulsed component of PSR B1706-44 should be detectable near threshold, whereas the unidentified GeV EGRET sources should be detectable if the superexponential cutoff energy is larger than ~ 30 GeV for relatively hard pulsar photon spectra ($\sim E^{-1.5}$).

A Introduction

Whereas gamma-ray pulsars are known to count amongst the brightest sources in the 1 - 30 GeV range (Lamb & Macomb, 1997), only their plerions, supernova shells and extragalactic sources appear to be visible at TeV energies. This is not unexpected, since pulsed γ -rays are known to be created in strong magnetic fields and magnetic pair production results in a superexponential cutoff above a characteristic energy E_o . Simulations of pair cascades have shown that E_o is usually in the 10 GeV region (Daugherty & Harding, 1996), but this cutoff depends on several parameters, such as the altitude of γ -ray production above the polar cap, the observer's viewing angle relative to the spin axis, and the magnetic inclination angle relative to the spin axis. A γ -ray/ $e^+/-$ cascade develops and those photons which escape pair creation (with $E < E_o$) are observable, resulting in a pulsed spectrum which is typically harder than E^{-2} for $E < E_o$. Based on this consideration, Nel & de Jager (1995) modelled the high energy γ -ray pulsed spectra of pulsars as

$$dN_\gamma/dE = k(E/E_o)^{-g} \exp(-(E/E_o)^b). \quad (1)$$

Whereas pulsar photon spectral indices between $g = 1.4$ and 2.1 are observed, harder spectra are theoretically possible (A.K. Harding, 2000, personal communication to O.C. de Jager). The constant k represents the monochromatic flux at the normalising energy $E_n \ll E_o$. We will normalise spectra at $E_n = 1$ GeV.

In the case of the outergap model for pulsars (Cheng, Ho, & Ruderman 1996), γ -ray production is expected to occur near the pulsar light cylinder, and the cutoff is expected to result from energetics arguments, rather than from magnetic pair production. In this case a larger E_o may be observable. Ground-based TeV γ -ray observations however provide firm upper limits on E_o . (Nel et al. (1993) gave a detailed discussion on this topic; see also Catanese & Weekes 1999.)

B Gamma-Ray Pulsar Spectral Parameters above 1 GeV

Table 1 shows the parameter results of a fit to the total pulsed spectra of the six brightest EGRET γ -ray pulsars. These spectral parameters reproduce the EGRET flux up to 30 GeV, and are consistent with the TeV pulsed limits. They also reproduce the GeV source catalog flux (Lamb & Macomb 1997). In the case of Vela and Geminga the cutoffs are well defined by the EGRET data and the errors on E_o are relatively small ($\sim 20\%$). In the case of Crab and PSR B1055-52, some evidence of a turnover is seen in the spectra above 10 GeV, although it is difficult to obtain reliable measures of E_o and b . In the case of PSR B1951+32 and PSR B1706-44 we see no evidence of a turnover up to 30 GeV, and a minimum value of $E_o = 40$ GeV (consistent with EGRET) was selected. This value is conservative with respect to the H.E.S.S. response. For those cases where E_o is not well defined, we have selected $b = 2$ (a value typical for a spectrum attenuated by magnetic pair production) to give conservative H.E.S.S. rates.

Using the H.E.S.S. collection area vs. energy $A(E)$ for any 2-telescope triggers (Konopelko 2000), we were able to calculate the expected rates R_p for pulsed γ -rays by integrating the product of $A(E)dN_\gamma/dE$ over all energies. The results for the six EGRET pulsars are shown in Table 1 (indicated by “ R_p ”). It is clear that the rate for PSR B1706-44 is the largest of all pulsars if E_o is not smaller than 40 GeV.

C H.E.S.S. Sensitivity for Pulsed γ -Ray Mission

It was shown by de Jager, Swanepoel & Raubenheimer (1987) and de Jager (1994) that the basic scaling parameter for any test for uniformity on the circle (given a test period) is given by $x = p\sqrt{n}$, where $p = R_p/(R_b + R_p)$ is the pulsed fraction, with R_p the pulsed rate and R_b the background rate. The total number of events is given by $N = (R_p + R_b)T$, with T the observation time. In this case the test statistic for uniformity for the general Beran (1969) class of tests is given by $B = x^2\Phi_B + c$, where Φ_B is derived from the intrinsic pulse profile, and c is the noise term. It was shown by Thompson (2000) that the pulse profiles above 5 GeV consist mostly of a single narrow peak, and it can be shown that $\Phi_B = 5.8$ for a 5%

TABLE 1. Gamma-ray spectral parameters above 1 GeV and corresponding H.E.S.S. rates and observation time for detection. Spectral references from Macomb & Gehrels (1999).

Object	k ($\times 10^{-8}$) ($\text{cm}^{-2}\text{s}^{-1}\text{GeV}^{-1}$)	g	E_o (GeV)	b	$F(> 1 \text{ GeV})$ ($\text{cm}^{-2}\text{s}^{-1}$)	R_p (hour^{-1})	T (10-hour days)
Crab	24.0	2.08	30	2	22	100	3
Vela	138	1.62	8.0	1.7	148	8	400
Geminga	73.0	1.42	5.0	2.2	76	$\ll 1$	-
PSR B1951+32	3.80	1.74	40	2	4.9	180	1
PSR B1055-52	4.00	1.80	20	2	4.5	8	420
PSR B1706-44	20.5	2.10	40	2	20	240	1

FWHM (single peak), if B is taken as the Z_m^2 test statistic with $m = 10$ harmonics (see e.g. de Jager, Swanepoel & Raubenheimer 1987). In this case $c = 20$.

A value of $x = 3$ would introduce a $\sim 3\sigma$ DC excess in a spatial analysis, but assuming that we have no imaging capability for E_o near the detection threshold, we have to rely on a timing analysis, which would give $Z_{10}^2 \sim 73$, or a chance probability of 7×10^{-8} if the period is known, but 0.03 after multiplying with the number of trials for a 6 hour observation if searching for periods as short as 50 ms. A confirming run (e.g. on a second night) should always be made to see if one of the few most significant periods from the previous run have repeated itself - in this case at the $\sim 10^{-7}$ level.

Using an additional topological software trigger, and selecting events by image size and angular shape, we were able to reject $\sim 99.2\%$ of the triggered background events, while retaining 95% of the source events. From a total background rate of 1 kHz (Konopelko 2000), we get $R_b = 8$ Hz. This allows us to calculate detection sensitivities for periodicities:

From the GeV source catalogue, we find that the galactic unidentified EGRET source (some may be pulsars - Lamb & Macomb 1997) fluxes range from $F(> 1 \text{ GeV}) = 1$ to $25 \times 10^{-8} \text{ cm}^{-2}\text{s}^{-1}$. Figures 1 and 2 give the H.E.S.S. sensitivity for a wide range of possible pulsar photon spectral indices between 1 and 2, and requiring a marginal detection within $T = 3$ to 6 hours (assuming a minimum “DC significance” of $x = 3$): Figures 1 and 2 respectively show E_o and T vs k , with the latter within the EGRET range as discussed above. Table 1 also shows T calculated in the same way, but assuming the spectral parameters of individual pulsars.

D Conclusions

It is clear that H.E.S.S. can only detect pulsars if E_o exceeds ~ 30 GeV. Even weak EGRET sources may be detectable if the spectra are as hard as E^{-1} , provided that E_o exceeds the levels prescribed by Figure 1. PSR B1706-44 (for which E_o is known to be at least as large as ~ 40 GeV) should be a H.E.S.S. candidate and




FIGURE 1. Figure 1 (Left panel): Parameter space (E_o vs k) for the detection of unknown pulsars within one night with H.E.S.S. using a timing analysis approach, and assuming $x = 3$. The three curves represent (from bottom to top) photon spectral indices of 1, 1.5 and 2.0. The solid line is for 3 hours of continuous observation, whereas the dashed lines (for the same set of spectral indices) represent a six-hour run. **Figure 2** (right panel): The observation time required to detect a pulsar as a function of k for a photon index of 1.5 and E_o as shown (also for $x = 3$).

other similar pulsars (such as PSR B1951+32) may be similarly detectable within one night. If one cannot detect a clear signal within a single night, an exact timing solution would be required to do a coherent analysis over a long period of time.

Whereas we have addressed the conservative polar cap model, any outergap component is expected to give a large value for E_o (which is no challenge for H.E.S.S.), but k may be small for such pulsars. This will be treated in a separate paper.

References

- Beran, R.J. 1969, Ann. Math. Statist., 40, 1196.
- Catanese, M. & Weekes, T.C. 1999, PASP, 111(764), 1193.
- Cheng, K.S., Ho, C. & Ruderman, M.A. 1986, ApJ, 300, 500.
- Daugherty, J.K. & Harding, A.K. 1996, ApJ, 458, 278.
- de Jager, O.C., Swanepoel, J.W.H. & Raubenheimer, B.C. 1986, A&A, 170, 187.
- de Jager, O.C. 1994, ApJ, 436, 239.
- Konopelko, A. 2000, these proceedings.
- Lamb, R.C. & Macomb, D.J., 1997, ApJ, 488, 872
- Macomb, D.J. & Gehrels, N. 1999, ApJ Suppl, 120, 335.
- Nel, H.I. et al. 1993, ApJ, 418, 836.
- Nel, H.I. & de Jager, O.C. 1995, Astr. Space Science, 230, 299.
- Thompson, D.J. 2000, these proceedings.